

BRIEF NOTE

GROWTH OF *PISIDIUM CASERTANUM* (POLI) IN WEST CENTRAL OHIO¹ALBERT J. BURKY, DANIEL J. HORNBACH² and C. M. WAY,³ Department of Biology, University of Dayton, Dayton, OH 45469

OHIO J. SCI. 81(1): 41, 1981

Clams of the family Sphaeriidae are found in most freshwater habitats. They are hermaphroditic and ovoviviparous (Mackie 1978), brooding their young in marsupial sacs formed as outgrowths of the gill filaments. In the genera *Sphaerium* and *Musculium*, several distinct larval stages can be present within a single adult, whereas in *Pisidium* only a single ontogenetic stage is present in a given adult (Heard 1977). Consequently, individuals of *Sphaerium* and *Musculium* produce a number of broods over a short period of time while in *Pisidium* more time is presumably required between successive series of broods.

Several life-history studies of sphaeriid clams are known: *Musculium* (Thomas 1963, Gale 1977, Mackie *et al* 1976, Mackie 1979), *Sphaerium* (Avolizi 1976, Mackie 1979) and *Pisidium* (Heard 1965, Ladle and Baron 1969, Holopainen 1979, Meier-Brook 1970, Thut 1969, Mackie 1979). Only the studies of Mackie (1979) and Thut (1969) present a full seasonal pattern of growth for North American species of *Pisidium*.

Clarke (1973) states that *Pisidium casertanum* (Poli) is perhaps the most common species of *Pisidium* and is truly cosmopolitan, occurring throughout Eurasia, Africa, Australia, and South, Central and North America. Populations of these clams are reported in habitats ranging from ephemeral ponds to benthic zones of deep lakes. Our study describes

aspects of the life-cycle of *P. casertanum* in a spring-fed stream with a relatively small annual temperature range.

Clams were collected monthly during the winter and bimonthly or weekly during the summer of 1974 and less regularly during 1975 from the west branch of Cedar Run at Cedar Bog, near Urbana, Ohio (USGS) map quadrangle Urbana West, Champaign County, Ohio: 40° 03.42'N 83°47.98'W). At certain times (see fig. 1) physical and chemical characteristics of the stream were recorded, including temperature and oxygen concentration (YSI Model 54 oxygen meter), conductivity (YSI Model 33 conductivity meter) and hardness (EDTA method, American Public Health Association 1976).

Usually more than 50 clams were removed from the stream sediment with a sieve (mesh opening = 0.5 mm) and fixed in the field with 12% neutral formalin. Shell length (greatest anterior-posterior dimension) was measured to the nearest 0.1 mm on a stage micrometer under a dissecting microscope. Samples collected within one week of each other were combined for a mean date. Samples were separated according to generation (adult *vs* juvenile) by the visual examination of frequency distributions of shell lengths. Generation samples with overlapping size distributions were separated by plotting cumulative frequency distributions on probability paper (Harding 1949, Cassie 1950, 1954).

Monthly variation in temperature and dissolved oxygen showed that in 1974 water temperature varied from 5.8 °C to 15 °C since the stream is fed from an underground source (Richard 1974), while

¹Manuscript received 29 May 1979 and in revised form 25 July 1980 (#79-34).

²Present address: Department of Biology, University of Virginia, Charlottesville, VA 22901.

³Present address: Department of Zoology, Miami University, Oxford, OH 45056.




FIGURE 1. Dissolved oxygen and water and air temperature in relation to months for the west branch of Cedar Run.

as expected, air temperature had a much greater range (fig. 1). Gilbert and Hennen (1974) have shown that air temperature in the bog area fluctuates much less than that in the surrounding areas. The oxygen content, although decreasing in summer and fall, was relatively high throughout the year, ranging from about 65% to 98% saturation. Mean annual conductivity was 564 μ mhos (range: 460–650 μ mhos), total hardness was 410–564 mg/l as CaCO₃, Ca hardness was 310–414 mg/l as CaCO₃ and Mg hardness was 160–310 mg/l as CaCO₃. The pH was usually > 7.2 (6.7–8.0), indicating that the stream, like the rest of the bog, is alkaline and not acidic. Bogs are by definition acidic; this habitat is actually a fen and alkaline in its water characteristics.

Size distributions of *P. casertanum* samples studied in 1974 revealed that there was one generation born per year with recruitment during April–August (fig. 2). The 1974 newborns were produced by clams born in 1972. Only *P. casertanum* with shell lengths of at least 3.0 mm are expected to contain extramarsupial larvae (those larvae ready to be born, Mackie 1979); therefore, these

clams take about 24 months to reach shell lengths of 3.0 mm, have one period of reproductive contribution and have a potential maximum life span of 30 to 33 months.

The life-cycle pattern of *P. casertanum* from Cedar Run is the only report with a complete sequence of samples over a full year for a stream population. The only other report we found of a stream population of *P. casertanum* is by Heard (1965). Although Heard (1965) didn't follow this population over an entire year, he reports one reproductive period during spring-summer. Holopainen (1979) found a 3 year life span for a littoral population in Lake Pääjärvi, Finland, with a single period of summer reproduction. He also reports a 3 year life span with 2 periods of reproduction for a population in Lake Esrom, Denmark. That (1969) reports a one year life span with one period of reproduction in early summer in Lake Washington. Mackie (1979) has examined 2 populations from ephemeral habitats with one population showing a single annual life-cycle with one period of reproduction while the other has 2 generations per year with complete summer replacement.




FIGURE 2. Mean shell length (solid black dot) in relation to months for generation samples of *Pisidium casertanum* for 1974. Vertical bars indicate the range and rectangles represent \pm standard deviation. The 3 generations were born in April-August of 1972, 1973 and 1974 (indicated in open circles) for the largest to smallest size distributions respectively. The number of clams represented on each date is given at the bottom of the figure with the percentage of animals for generations indicated in or near the rectangles.

The life cycle of *P. casertanum* ranges from 2 generations per year with complete replacement (Mackie 1979) to the simple annual (Thut 1969, Mackie 1979) to three year life spans with one or two periods of reproduction (Holopainen 1979). *Pisidium casertanum* at Cedar Run (fig. 2) is well within this range with a maximum life span of 30 to 33 months with a single period of reproductive contribution after 24 months of growth and maturation.

Acknowledgments. We would like to thank the Ohio Historical Society for permission to collect clams at Cedar Bog and Dr. G. L. Mackie for having confirmed our identification of these clams as *Pisidium casertanum* (Poli). This study has been supported in part by grants to Albert J. Burky from the Ohio Biological Survey and the University of Dayton Research Council.

LITERATURE CITED

American Public Health Association 1976 Standard methods for the examination of water and wastewater (14th ed.) p. 1193.

Avolizi, R. J. 1976 Biomass turnover in populations of viviparous sphaeriid clams: Comparisons of growth, fecundity, mortality and biomass production. *Hydrobiologia* 51: 163-180.

Cassie, R. M. 1950 Analysis of polymodal frequency distribution by the probability paper method. *N. Z. Sci. Rev.* 8: 89-91.

——— 1954 Some uses of probability paper in the analysis of size frequency distributions. *Aust. J. Mar. Freshwater Res.* 5: 513-522.

Clarke, A. H. 1973 The freshwater molluscs of the Canadian Interior Basin. *Malacologia* 13: 1-509.

Gale, W. F. 1977 Growth of the fingernail clam, *Sphaerium transversum* (Say) in field and laboratory experiments. *Nautilus* 91: 8-12.

Gilbert, G. E. and R. W. Hennen 1974 Frequency and structure of temperature inversions in Cedar Bog. pp. 14-15. In: C. C. King and C. M. Frederick (eds.) Cedar Bog Symposium. *Ohio Biol. Surv. Inf. Cir.* 4: 71 p.

Harding, J. P. 1949 The use of probability paper for the graphical analysis of polymodal frequency distributions. *J. Marine Biol. Asso. U.K.* 28: 141-153.

- Heard, W. H. 1965 Comparative life histories of North American pill clams (Sphaeriidae: *Pisidium*). *Malacologia* 2: 381-411.
- 1977 Reproduction of fingernail clams (Sphaeriidae: *Sphaerium* and *Musculium*). *Malacologia* 16: 421-455.
- Holopainen, I. J. 1979 Population dynamics and production of *Pisidium* species (Bivalvia, Sphaeriidae) in the oligotrophic and mesohumic Lake Pääjärvi, Southern Finland. *Arch. Hydrobiol. Suppl.* 54: 466-508.
- Ladle, M. and F. Baron 1969 Studies on three species of *Pisidium* (Mollusca: Bivalvia) from a chalk stream. *J. Anim. Ecol.* 38: 407-413.
- Mackie, G. L. 1978 Are sphaeriid clams ovoviviparous or viviparous? *Nautilus* 92: 145-146.
- 1979 Growth dynamics in natural populations of Sphaeriid clams (*Sphaerium*, *Musculium*, *Pisidium*). *Canadian J. Zool.* 57: 441-457.
- , S. U. Qadri and A. H. Clarke 1976 Intraspecific variations in growth, birth periods, and longevity of *Musculium securius* (Bivalvia: Sphaeridae) near Ottawa, Canada. *Malacologia* 15: 433-446.
- Meier-Brook, V. C. 1970 Investigations on the biology of some *Pisidium* species (Mollusca; Eulamellibranchiata; Sphaeriidae). *Arch. Hydrobiol. Suppl.* 38: 73-150.
- Richard, B. H. 1974 The relationship of Cedar Bog to the ancestral Teays River System. pp. 3-7. *In*: C. C. King and C. M. Frederick (eds.) Cedar Bog Symposium. Ohio Biol. Surv. Inf. Cir. 4. 71 p.
- Thomas, G. J. 1963 Study of a population of sphaeriid clams in a temporary pond. *Nautilus* 77: 37-43.
- Thut, R. N. 1969 A study of the profundal bottom fauna of Lake Washington. *Ecol. Monogr.* 39: 79-100.