
PLANKTON POPULATION OF CERTAIN LAKES AND STREAMS IN THE ROCKY MOUNTAIN NATIONAL PARK, COLORADO

FLOYD J. BRINLEY,

University of Toledo, Toledo, Ohio

An extensive survey of the plankton population of the streams in the Ohio River watershed was made from 1939-1942 (Brinley and Katzin, 1944). This study revealed the fact that the northern tributaries of the Ohio River support a much larger phytoplankton population in species and numbers than the southern tributaries. This difference in plankton distribution is attributed to the fact that the northern streams flow through densely populated areas, the farm lands are fertile and the surface waters are generally polluted with human and industrial wastes. The southern streams, on the other hand, pass through sparsely populated regions where the soils are poor and the streams are less likely to be enriched with organic pollution. The algae population seems to be directly correlated with the amount of decomposed organic matter in the stream. Studies of the White (Brinley, 1942a) and the Cumberland Rivers (Brinley, 1942b) clearly showed that isolated sources of heavy organic pollution greatly increased the growth of algae, in numbers and species, in the stream below the entrance of the sewage.

It is desirable to obtain more information on the relation of stream fertility to the plankton population by studying streams which are free from organic pollution. Such streams do not exist in the state of Ohio so it seemed advisable to study the problem in sparsely populated mountain districts where the streams are free from human wastes and where the decomposition of natural organic matter, vegetation, leaves, humus, etc., is at a minimum.

Permission was freely given by the National Park Service to conduct these studies in the Rocky Mountain National Park. The writer wishes to express his appreciation to David Canfield, Superintendent, J. Barton Herschler, Chief Ranger, and Edwin C. Alberts, Park Naturalist, for all facilities of the Park which were so graciously given.

METHODS AND PROCEDURE

The present studies were made in the Park from June 13 to July 22, 1949. In the beginning of the study, samples of water were collected in wide mouth bottles and an attempt was made to make qualitative and quantitative determinations of the population of plankton and to express the quantitative results in parts per million as recorded in the previous publications (Brinley, 1942). However, it was soon found that the plankton population of these streams and lakes was so low

that collections of small samples yielded very few individuals and the volume was less than one part per million as compared with several parts per million as found in the Ohio River Basin. It was decided, therefore, to discontinue the small sample collections and to use a plankton net. Approximately ten to twenty gallons of water were filtered through the net by sweeping the net through quiet water or allowing running water to flow through the net. The fifteen milliliters of catch were then concentrated to one ml. by centrifuging. This method was largely qualitative and thus no volume determinations were made. The lake samples were taken from the shore line on the leeward side of the lake. Samples from the streams were largely taken from boulders along the shore by allowing the water to flow through the net.




FIG. 1. Map of the area from which collections were made.

The stream beds and lakes in the Park were glacier formed and thereby in many cases two to four lakes are terraced up the gorges and during high water these lakes are connected by a flowing stream. The lakes studied were conveniently arranged in related groups as they occur in the common valley (see map tracing). In most cases the lakes had stony bottoms and were partly filled with loose boulders that had descended from rocky ledges above. The rocks along the shore line were covered with unidentified filamentous algae and a layer of silt of various depths covered the bottom between the rocks and boulders. During the

month of June, snow banks extended to the edge of the water in most of the lakes of high elevations.

The typical mountain streams flowed over rocky valley floors and during high water in June the samples collected contained large amounts of sand which seriously interfered with the plankton determinations. Unidentified filamentous algae covered the rocks.

LAKES

DISTRIBUTION OF PLANKTON

GROUP I

Lake	Elevation Ft.	Temp. C.	Plankton	Relative Population	Date				
Bear	9,600	10	DIATOMS						
			<i>Asterionella gracillima</i>	abundant	6/18				
			<i>Diatoma vulgare</i>	few					
			<i>Navicula</i> sp.....	few					
			<i>Cymbella</i> sp.....	few					
			<i>Synedra</i>	scarce					
			DESMIDS						
			<i>Closterium subcostatum</i>	scarce					
			<i>Staurastrum</i> sp.....	scarce					
			GREEN ALGAE						
			<i>Cryptomonas ovalis</i>	scarce					
			<i>Volvox</i>	scarce					
			PROTOZOA						
			<i>Peridinium</i>	scarce					
			COPEPODS						
<i>Cyclops</i>	few								
Nymph	9,800	16	DIATOMS		7/15				
			<i>Asterionella gracillima</i>	abundant					
			<i>Diatoma vulgare</i>	few					
			<i>Navicula</i> sp.....	few					
			DESMIDS						
			<i>Closterium</i> sp.....	few					
			GREEN ALGAE						
			<i>Dinobryon</i> sp.....	abundant					
			<i>Volvox</i>	one colony					
			<i>Cryptomonas ovalis</i>	scarce					
			CLADOCERA						
			<i>Daphnia pulex</i>	abundant					
			Dream	10,000	17	DIATOMS		6/30	
						<i>Navicula</i> sp.			
						<i>Diatoma vulgare</i>	abundant		
DESMIDS									
<i>Penium closterioides</i>	few								
<i>Micrasterias radiosa</i>	few								
<i>Staurastrum</i> sp.....	few								
COPEPODS									
<i>Cyclops</i>	few								
ROTIFERS									
<i>Anuraea</i>	abundant								
Dream	10,000	7				DIATOMS		6/30	
						<i>Asterionella</i>	abundant		
						<i>Diatoma vulgare</i>	few		
						<i>Synedra</i>	few		
			GREEN ALGAE						
			<i>Cryptomonas erosa</i>	scarce					
			Dream	10,000	8	DIATOMS		7/8	
						<i>Asterionella</i>	abundant		
						<i>Diatoma vulgare</i>	few		
						DESMIDS			
						<i>Microsterias radiosa</i>	few		
						COPEPODS			
						<i>Cyclops</i>	scarce		
						ROTIFERS			
						<i>Anuraea</i>	scarce		

DISTRIBUTION OF PLANKTON—(Continued)

LAKES

GROUP I—Continued

Lake	Elevation Ft.	Temp. C.	Plankton	Relative Population	Date
Emerald	10,200	8	DIATOMS		
			<i>Asterionella</i>	abundant	
			<i>Diatoma vulgare</i>	few	
			DESMIDS		
			<i>Micrasterias radiosa</i>	few	
			GREEN ALGAE		
			<i>Cryptomonas erosa</i>	scarce	

GROUP II

The Loch	10,700	8	DIATOMS		
			<i>Diatoma vulgare</i>	few	
			GREEN ALGAE		
			<i>Volvox</i>	abundant	
			<i>Cryptomonas erosa</i>	few	
			COPEPODS		
			<i>Cyclops</i>	few	
Lake-of-Glass	10,800	6	DIATOMS		
			<i>Asterionella</i>	abundant	
			GREEN ALGAE		
			<i>Volvox</i>	abundant	
			<i>Cryptomonas erosa</i>	few	
			COPEPODS		
			<i>Cyclops</i>	few	
			ROTIFERS		
			<i>Noteus</i>	few	
Sky-Pond	11,100	6	DIATOMS		
			<i>Asterionella</i>	few	
			GREEN ALGAE		
			<i>Volvox</i>	abundant	
			<i>Cryptomonas erosa</i>	few	

GROUP III

Mills	11,500	9	DIATOMS		
			<i>Navicula</i>	few	
			DESMID		
			<i>Cosmarium</i>	few	
			ROTIFERS		
			<i>Anuraea</i>	few	
Jewel	11,600	9	DIATOMS		
			<i>Navicula</i> sp.		
			<i>Diatoma vulgare</i>		

GROUP IV

Fern	9,500	7	DIATOMS		
			<i>Navicula</i> sp.....	abundant	
			<i>Diatoma vulgare</i>	many	
			<i>Synedra</i>	few	
			GREEN ALGAE		
			<i>Cryptomonas erosa</i>	few	
Odessa	10,000	5	DIATOMS		
			<i>Navicula</i> sp.....	few	
			ROTIFERS		
			<i>Anuraea</i>	few	

GROUP V (Miscellaneous)

Bierstadt	9,500	13	DIATOMS		
			<i>Diatoma vulgare</i>	many	
			<i>Navicula</i>		
			DESMIDS		
			<i>Cosmarium constrictum</i>	few	
			<i>Closterium subcostatum</i>	few	
			GREEN ALGAE		
			<i>Euglena viridis</i>	few	
			<i>Phacus pyrum</i>	very few	
			PROTOZOA		
<i>Vorticella</i>	very few				

DISTRIBUTION OF PLANKTON—(Continued)

LAKES

GROUP V—Continued

Lake	Elevation Ft.	Temp. C.	Plankton	Relative Population	Date
Cub	8,600	17	DIATOMS <i>Diatoma vulgare</i>	few	6/21
			CLADOCERA <i>Daphnia pulex</i>	very abundant	
			ROTIFERS <i>Anuraea</i>	abundant	
Chipmunk	9,900	17	DIATOMS <i>Diatoma vulgare</i> <i>Navicula</i> sp.....	few few	6/23
			DESMIDS <i>Euastrum abruptum</i> <i>Cosmarium constrictum</i> <i>Closterium subcostatum</i>	few few few	
			CLADOCERA <i>Daphnia pulex</i>	few	
			ROTIFERS <i>Anuraea</i> <i>Polyarthra</i>	abundant few	
Lily	9,900	18	DIATOMS <i>Navicula</i>	abundant	6/24
			DESMIDS <i>Micrasterias radiosa</i>	few	
			GREEN ALGAE <i>Chlamydomonas</i> sp..... <i>Volvox</i> <i>Euglena</i> sp.....	few few very few	
			PROTOZOA <i>Ceratium</i>	very few	
			CLADOCERA <i>Daphnia pulex</i>	many	
			COPEPODS <i>Cyclops</i>	few	
			ROTIFERS <i>Anuraea</i> <i>Noteus</i>	few few	
Haiyaha	10,700	6	DIATOMS <i>Asterionella</i> <i>Diatoma vulgare</i>	few few	7/8
			GREEN ALGAE <i>Cryptomonas ovata</i>	very few	
			ROTIFERS <i>Anuraea</i> <i>Noteus</i>		
			COPEPODS <i>Cyclops</i>	few	

STREAMS

Streams	Location	Temp. C.	Plankton	Relative Population	Date
Big Thompson	Below "Pool"	6	DIATOMS <i>Navicula</i> <i>Synedra acus</i>	many	6/18
			DESMIDS <i>Closterium subcostatum</i>		
		8	DIATOMS <i>Navicula</i> <i>Nitzschia</i>	many few	7/10
		9	DIATOMS <i>Navicula</i>	few	7/20

DISTRIBUTION OF PLANKTON—(Continued)

STREAMS

Stream	Location	Temp. C.	Plankton	Relative Population	Date
	Stead's Hotel	8	DIATOMS		7/3
			<i>Navicula</i> sp.....	few	
			<i>Diatoma vulgare</i>	few	
			DESMIDS		
			<i>Closterium</i> sp.....	few	
			ROTIFERS		
			<i>Anuraea</i>		
		8	DIATOMS		7/10
			<i>Navicula</i>	few	
		11	<i>Navicula</i>	few	7/20
	Moraine Park	8	DIATOMS.....		6/27
			<i>Navicula</i>	few	
			<i>Diatoma vulgare</i>	few	
			DESMIDS		
			<i>Cosmarium constrictum</i>	few	
			<i>Closterium subcostatum</i>		
			PROTOZOA		
			<i>Loxodes</i>		
		8	DIATOMS		7/10
			<i>Navicula</i> sp.....	few	
			<i>Diatoma vulgare</i>	few	
			<i>Closterium subcostatum</i>	few	
			<i>Cosmarium constrictum</i>	few	
		11	DIATOMS		7/20
			<i>Diatoma vulgare</i>	few	
			<i>Navicula</i> sp.		
			DESMIDS		
			<i>Cosmarium constrictum</i>		
			<i>Closterium subcostatum</i>		
Glacier Creek	Below Bear Lake	8	DIATOMS		7/2
			<i>Asterionella</i>	many	
			<i>Navicula</i>	many	
			<i>Nitzschia</i>	few	
			<i>Synedra</i>	few	
			GREEN ALGAE		
			<i>Cryptomonas erosa</i>	very few	
	Sprague's Lodge	8	DIATOMS		7/2
			<i>Asterionella</i>	many	
			<i>Navicula</i>	few	
			<i>Nitzschia</i>	few	
			<i>Diatoma vulgare</i>	few	
			DESMIDS		
			<i>Closterium</i>	few	
	Mouth of Mill Creek	8	DIATOMS.....		7/2
			<i>Asterionella</i>	few	
			<i>Navicula</i>	few	
			<i>Nitzschia</i>	few	
			GREEN ALGAE		
			<i>Cryptomonas erosa</i>	very few	
		10	DIATOMS		7/15
			<i>Asterionella</i>	few	
			<i>Navicula</i>	few	
			DESMIDS		
			<i>Micrasterias radiosa</i>	few	
Fall River	Chasm Falls	5	DIATOMS		7/9
			<i>Navicula</i>	very few	
		8	DIATOMS		7/16
			<i>Navicula</i>	few	
	End-of-Valley Camp	8	DIATOMS		7/9
			<i>Navicula</i> sp.....	few	
			<i>Diatoma vulgare</i>	few	
		9	DIATOMS		7/16
			<i>Navicula</i> sp.....	few	

DISTRIBUTION OF PLANKTON—(Continued)

STREAMS

Stream	Location	Temp. C.	Plankton	Relative Population	Date
	Fall River Lodge	5	DIATOMS <i>Navicula</i> sp.....	few	7/6
			<i>Diatoma vulgare</i>	few	
			ROTIFERS <i>Noteus</i>	very few	
			PROTOZOA <i>Actinosphaerium sol</i>	very few	
		8	DIATOMS <i>Navicula</i> sp.....	few	7/9
			<i>Diatoma vulgare</i>	few	
			DESMIDS <i>Closterium</i> sp.....	few	
			PROTOZOA <i>Cyclidium</i>		
	Highway Bridge U. S. 34	7	DIATOMS <i>Navicula</i>	few	6/28
			<i>Diatoma vulgare</i>	few	
		8	DIATOMS <i>Navicula</i> sp.....	very few	7/9
			PROTOZOA <i>Cyclidium</i>	very few	
	Below Sheep Lake	7	DIATOMS <i>Navicula</i>	few	6/28
			GREEN ALGAE <i>Tetraspora</i>	few	
	Above Estes Park		DIATOMS <i>Navicula</i>	few	6/28
			<i>Diatoma vulgare</i>	few	
			DESMIDS <i>Cosmarium rostratum</i>	few	

DISCUSSION AND SUMMARY

A study of the data presented shows that the diatoms were the most abundant group of planktons in both the lakes and streams, followed in numbers by the desmids and an occasional green flagellate and protozoan. In some lakes *Daphnia*, *Cyclops* and rotifers were abundant. The small number of species may in a large part be due to the low temperature, in most cases below 10 degrees Centigrade. The species found in the Park as listed in this paper are the typical cold water forms that occur in the Ohio Basin (Brinley and Katzin 1942). The total plankton volume in the mountain streams is much lower than the volume of the same species in the Ohio River Basin under similar temperature conditions, which I believe can be attributed to the fertilizing action of the organic pollutants in the Ohio stream.

A comparative study of the streams and lakes in the Park showed no specific differences in the plankton algae, however, the population density of individual species were generally higher in the lakes. *Daphnia* and *Cyclops* were abundant in Bear, Cub and Lily lakes. Rotifers were numerous in Chipmunk, Cub, and Lily lakes. Hellgrammites and leeches were abundant in the latter lakes and these lakes also supported a heavy population of water lilies.

A careful study of the distribution records indicates a possible similarity in the plankton algae in closely related lakes. The lakes in group one, Bear, Nymph, Dream and Emerald are all located in the Tyndall Glacier Gorge. The characteristic specie of algae found in three of these lakes is the diatom, *Asterionella*. It is also interesting to note that this diatom was widely distributed in Glacier Creek which receives the outlet from Bear Lake and Glacier Gorge. The individual specimens of this fragile diatom in Glacier Creek were always broken which indi-

cates that the origin of this specie was in the lakes and not a direct product of the stream. A few *Asterionella* were, however, found in Lake Haiyaha, Lake-of-Glass, Sky Pond, and The Loch. Volvox was found abundantly in the Loch Vale Lakes (group 2); The Loch, Lake-of-Glass and Sky Pond, and only occasionally elsewhere.

The stream plankton was typically diatoms. There is a tendency for the desmids to increase in the Big Thompson River as it flows through the meadows in Moraine Park.

REFERENCES

- Brinley, F. J., and Katzin, L. J.** 1944. Report of the U. S. Public Health Service, Ohio River Pollution Control. Suppl. to Part 2, pp. 1279-1368.
- Brinley, F. J.** 1942 (a). The effect of pollution upon the plankton population of the White River, Indiana. Investigation of Indiana Lakes and Streams, 2, 137-143. Indiana Department of Conservation.
- Brinley, F. J.** 1942 (b). The effect of sewage from Nashville upon the plankton population of the Cumberland River. Jour. of the Tenn. Acad. of Science 17: 179-183.
- Brinley, F. J., and Katzin, L. J.** 1942. Distribution of stream plankton in the Ohio River System. Am. Mid. Naturalist 27, No. 1, 173-182.
-